

Outcomes following Prenatal Exposure to Dolutegravir: the Dolomite-EPPICC study

C. Thorne¹, V. Rasi¹, K. Aebi-Popp², L. Ene³, M. Florida⁴,
N. Mendoza-Palomar⁵, L. Prieto⁶, L. Ragone⁷, R. Sconza¹, C. Giaquinto⁸,
V. Vannappagari⁷ for the Dolomite-EPPICC study group

1. UCL Great Ormond Street Institute of Child Health, London, UK; 2. Bern University Hospital, Switzerland; 3. Victor Babes Hospital, Bucharest, Romania; 4. Istituto Superiore di Sanità, Rome, Italy; 5. Hospital Universitari Vall d'Hebron, Barcelona, Spain; 6. Hospital Universitario 12 de Octubre, Madrid, Spain; 7. ViiV Healthcare, Research Triangle Park, USA; 8. Fondazione Penta Onlus, Padua, Italy

Disclosure slide

- This study has been supported by ViiV Healthcare



Outside the current work, the presenting author has received support from AbbVie and ViiV, plus Public Health England, UK Medical Research Council, European Commission, PENTA Foundation.

Background

- Dolutegravir (DTG) is an integrase strand inhibitor approved for treatment of HIV in adults and adolescents since 2013
- In 2018, **Tsepamo Study, Botswana** reported a significantly increased neural tube defect (NTD) risk in women conceiving on DTG (0.94%)¹, leading to a safety alert.
- Updated analysis of NTD prevalence (August 2014 to March 2019)²
 - 5 NTDs among 1,683 deliveries (**0.30%**, 95% CI 0.13-0.69) in women on DTG at conception
 - **0.10%** 95% CI 0.06-0.17 in women on non-DTG ART at conception
 - **0.08%**, 95% CI 0.06-0.10 in HIV-uninfected women
 - prevalence difference for DTG vs. non-DTG at conception remained statistically significant
- **Antiretroviral Pregnancy Registry** reported 1 NTD with 312 periconception DTG exposures (NTD prevalence of 0.3% for DTG; overall risk of defects with periconception DTG was 3.2% (data to July 2019)³

¹Zash et al, NEJM 2018; ²Zash et al, NEJM 2019; ³Vannappagari et al EACS 2019

Aim

- To assess pregnancy and neonatal outcomes following DTG use during pregnancy in real-world European settings

Our objectives were to describe:

- the characteristics of pregnant women receiving DTG-based regimens
- the frequency of adverse pregnancy and birth outcomes, by earliest timing of DTG exposure


Methods

- The **Dolomite Study** was set up in 2017 to address the use and safety of DTG in pregnant women and exposed infants in Europe and Canada and is conducted within the NEAT-ID network and EPPICC (the European Pregnancy and Paediatric Infections Cohort Collaboration)
- **Dolomite-EPPICC** involves pooled analysis of observational data on DTG-exposed pregnancies from participating studies, following periodic data mergers
- Pseudonymised individual patient data collected from participating studies using a data specification based on a modified HIV Data Exchange Protocol (www.hicdep.org)
- Analysis of prospectively collected individual patient data (i.e. ARV exposure data collected before outcome is known)
- Data merger included
 - All pregnancies with any prenatal DTG exposure
 - With birth outcomes reported by Feb 2019

Definitions

Periconception exposure	Initial exposure within the first 6 weeks of estimated gestational age
Later 1 st trimester (T1)	Initial exposure started later in T1 (after 6 weeks estimated gestational age)
2 nd / 3 rd trimester (T2/T3)	Initial exposure started after 12 weeks estimated gestational age
Induced abortion	Voluntary termination of pregnancy before 22 weeks estimated gestational age
Spontaneous abortion	Death of a fetus or expulsion of the products of conception before 22 weeks gestational age
Low birth weight	Birth weight of <2500 grams
Very low birth weight	Birth weight of <1500 grams
Preterm birth	Birth of live infant at <37 completed weeks gestation
Stillbirth	Death of a fetus occurring at 22 weeks of gestation or more, or for situations in which the gestational age is unavailable, a fetus weighing at least 500 grams

Results



Pregnancy / maternal characteristics

- 453 pregnancies in 428 women
- Pregnancies reported from 6 countries
 - 347 (76.6%) UK and Ireland
 - 45 (9.9%) Spain
 - 29 (6.4%) Switzerland
 - 29 (6.4%) Italy
 - 3 (0.7%) Romania
- Timing of earliest DTG exposure in 453 pregnancies
 - periconception: 317 (70.0%)
 - later in 1st trimester: 31 (6.8%)
 - 2nd or 3rd trimester: 105 (23.2%)

	428 women	N (%)
Ethnicity N=428	Black African	229 (53.5)
	Black other	35 (8.2)
	White	129 (30.1)
	Other	35 (8.2)
Mode of HIV acquisition N=394	Heterosexual	326 (82.7)
	Injecting drug use	11 (2.8)
	Vertical	42 (10.6)
	Other	15 (3.8)
History of AIDS N=353	Yes	30 (8.5)
HCV status N=395	HCV seropositive	24 (6.1)
HBV status N=393	HBsAg positive	14 (3.6)

Outcomes, by earliest DTG exposure

	Total DTG exposed	Earliest exposure to DTG		
		Periconception	Later T1	T2/T3
Total outcomes, N	463*	325	31	106
Live births	417 (90.1%)	280 (86.1%)	30 (96.8%)	106 (100%)
Stillbirths	5 (1.1%)	5 (1.5%)	0	0
Spontaneous abortions	23 (5.0%)	23 (7.2%)	0	0
Induced abortions	18 (3.7%)	17 (5.2%)	1 (3.2%)	0

*includes outcomes from 10 twin pregnancies

Neonatal Outcomes: 400 singleton, live births

	Total DTG exposed	Earliest exposure to DTG		
		Periconception	Later T1	T2/T3
Total, N	400	266	30	104
Gestational age				
<34 weeks	12 (3.1%)	8 (3.0%)	1 (3.3%)	3 (2.8%)
34-36 weeks	39 (9.7%)	24 (9.0%)	2 (6.7%)	13 (12.5%)
≥ 37 weeks	334 (83.5%)	222 (83.5%)	26 (86.7%)	86 (82.7%)
missing	15 (3.7%)	12 (4.5%)	1 (3.3%)	2 (1.9%)
Birth weight				
<1500g	12 (3.0%)	8 (3.1%)	1 (3.3%)	3 (2.9%)
1500-2499g	36 (9.0%)	23 (8.6%)	2 (6.7%)	11 (10.6%)
≥2500g	342 (85.5%)	230 (86.5%)	26 (86.7%)	86 (82.7%)
missing	10 (2.5%)	5 (1.8%)	1 (3.3%)	4 (3.8%)

Birth defects

- Among the 417 live-born infants there were 17 with reported birth defects (4.1%, 95% CI 2.4, 6.5); one infant had 2 defects.

% of infants with birth defects by timing of earliest exposure to DTG:

Periconception	12/266	4.5%	(95% CI 3.9, 5.1)
Later T1	1/30	3.3%	(95% CI 0.08, 17.2)
T2/T3	4/104	3.8%	(95% CI 1.1, 9.6)

- No defects in stillborn infants
- Of 18 outcomes of induced abortion, there was 1 carried out due to identified birth defects
 - at gestation 29 weeks for neuronal migration disorder and severe microcephaly
 - periconception DTG exposure

Birth defects: details

Organ system	Exposure	Birth defect
Heart N=3	PC	Patent Foramen Ovale
	PC	Interatrial communication – ostium secundum
	PC	Septal defect
Genitourinary N=7	PC	2 x Congenital hydronephrosis
	PC	Ectopic Kidney
	PC	3 x Hypospadias*
	T2/3	Hypospadias
Gastrointestinal N=2	T2/3	Duodenal atresia and stenosis
	PC	Gastroschisis
Limb N=3	PC	2 x Polydactyly*
	T2/3	Polydactyly
Other N=3	Later T1	Ankyloglossia
	T2/3	Hyperpigmentation on back
	PC	Naevus flammeus

*1 infant had hypospadias and polydactyly

PC= periconception

Birth defects: details

Organ system	Exposure	Birth defect	EuroCAT?
Heart N=3	PC	Patent Foramen Ovale	No
	PC	Interatrial communication – ostium secundum	Yes
	PC	Septal defect	Yes
Genitourinary N=7	PC	2 x Congenital hydronephrosis	Yes
	PC	Ectopic Kidney	Yes
	PC	3 x Hypospadias*	Yes
	T2/3	Hypospadias	Yes
Gastrointestinal N=2	T2/3	Duodenal atresia and stenosis	Yes
	PC	Gastroschisis	Yes
Limb N=3	PC	2 x Polydactyly*	Yes
	T2/3	Polydactyly	Yes
Other N=3	Later T1	Ankyloglossia	No
	T2/3	Hyperpigmentation on back	No
	PC	Naevus flammeus	No

*1 infant had hypospadias and polydactyly

PC= periconception

Perinatal mortality

Stillbirths

- 5 Stillbirths: 4 female, 1 unknown sex; 4 preterm, 1 term
- All exposed to periconception DTG
- None with birth defects reported
- Stillbirth rate: 11.8 per 1000 (95% CI 3.9, 27.4)

Neonatal death

- 1 death in an extremely preterm infant
 - born at 23 weeks gestation
- Died on 2nd day of life
- Periconception DTG exposure

Perinatal mortality rate: 14.2 per 1000 (95% CI 5.2, 30.7)

Comparative data from National Surveillance of HIV in Pregnancy and Childhood (England & Wales), 2010-2016

	Rate /1000	95% CI
Stillbirth	7.4	(5.7, 9.5)
Neonatal death	4.2	(2.8, 5.7)
Extended perinatal mortality	11.5	(9.3, 14.0)

Conclusions

- This is the largest study to date of DTG use in pregnancy in Europe in which 70% of 453 pregnancies had periconception DTG exposure
- Overall, 4.1% prevalence of birth defects (3.1% if considering only EuroCAT defects)
- No NTDs were reported – but:
 - NTD are rare events ($\approx 0.1\%$ birth prevalence)
 - sample size of <300 pregnancies with periconception DTG exposure
 - cannot rule out or confirm any potential association of DTG with NTD
 - 2000 exposures needed to rule out a 3-fold increase for NTDs
- The birth defect rate and pattern add to the current evidence base on periconception DTG use and safety
- This study is ongoing, in order to provide additional data from European settings

Acknowledgements

Many thanks to the study participants in the following cohorts and studies:

- Italian Group on Surveillance of Antiretroviral Treatment in Pregnancy
- NENEXP Study (Catalonia)
- Swiss Mother and Child HIV Cohort Study (MoCHiV)
- UK / Ireland National Surveillance of HIV in Pregnancy and Childhood (NSHPC)
- Madrid Cohort of HIV-Infected Mother-Infant Pairs
- Victor Babes Hospital Cohort
- This study was supported by ViiV Healthcare

